

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
International General Certificate of Secondary Education

**MARK SCHEME for the May/June 2009 question paper  
for the guidance of teachers**

**0580, 0581 MATHEMATICS**

**0580/04, 0581/04** Paper 4 (Extended), maximum raw mark 130

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



<b>Page 2</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – May/June 2009</b>	<b>0580, 0581</b>	<b>04</b>

### Abbreviations

cao	correct answer only
cso	correct solution only
dep	dependent
ft	follow through after error
isw	ignore subsequent working
oe	or equivalent
SC	Special Case
www	without wrong working

<b>1 (a)</b>	<b>(\$ 450)</b>	<b>B2</b>	<b>M1</b> for $650 \div (9 + 4) \times 9$ ( $\div 14$ does not imply $9 + 4$ )
<b>(b) (i)</b>	<b>(\$ 120)</b>	<b>B2</b>	<b>M1</b> for $0.8 \times 150$ o.e.
<b>(ii)</b>	<b>(\$ 80 ft)</b>	<b>B2 ft</b>	<b>M1</b> for $(150 - \text{their(b)(i)}) \div 0.375$ o.e. only if +ve. After <b>M0, SC1</b> for answer 320
<b>(c) (i)</b>	<b>(\$ 441)</b>	<b>B2</b>	<b>M1</b> for $400 \times 1.05^2$ o.e. or for answer 41
<b>(ii)</b>	$\frac{1}{2}$ their <b>(i)</b> $\div 400 \times 100$ o.e.  <b>5.125</b> or <b>5.13</b> or <b>5.12</b> c.a.o. www3	<b>M2</b>  <b>A1</b>	<b>If use Simple Int in (i), M0, M0 in this part</b> i.e. a full explicit method for $r$ If <b>M0</b> , <b>M1</b> for $\frac{400 \times r \times 2}{100} = \text{their (i)} - 400$  or their <b>(i)</b> $\div 400 \times 100$ then $- 100$  or $\frac{\text{their (i)} - 400}{400} \times 100$ (s.o.i. by 10.25)  If still <b>M0, SC1</b> for answers 55.125 or 55.12 or 55.13 or 55.1 or 0.05125 or 0.0512 or 0.0513  <b>[11]</b>

<b>2 (a)</b>	<b>1</b>	<b>B1</b>	
<b>(b)</b>	<b>2.5</b> o.e.	<b>B1</b>	
<b>(c)</b>	<b>2.96</b> c.a.o.	<b>B2</b>	If <b>B0, M1</b> for $15 \times 1 + 10 \times 2 + 7 \times 3 + 5 \times 4 + 6 \times 5 + 7 \times 6$ (allow one slip) implied by 148 seen Ignore subsequent rounding
<b>(d)</b>	$60 \times 2.95 (= 177)$ their 177 $-$ their 148 (or $50 \times$ their 2.96)  (Mean of new rolls $\Rightarrow$ ) <b>2.9</b> c.a.o. www3	<b>M1</b> <b>M1</b> <b>A1</b>	<b>Dependent</b> on first <b>M</b> and <u>only if</u> positive <b>or M1</b> for $\frac{\text{their } 148(50 \times \text{their } 2.96) + x(\text{or } 10x)}{60} = 2.95$  then <b>M1</b> for $x(\text{or } 10x) = 60 \times 2.95 - \text{their } 148$ (or $50 \times$ their 2.96) and <u>only if</u> positive  <b>[7]</b>

<b>Page 3</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – May/June 2009</b>	<b>0580, 0581</b>	<b>04</b>

<b>3 (a)</b>	$(\sin P) = \frac{48}{0.5 \times 10 \times 14}$ o.e. <u>fraction</u>  $P = 43.29\dots$ cao	<b>M2</b>  <b>A1</b>	<b>M1</b> for $0.5 \times 10 \times 14 \sin P = 48$ o.e. <b>Allow <math>0.5 \times 10 \times 14 \sin 43.3 = 48</math> for M1 but no further credit</b>
<b>(b)</b>	$10^2 + 14^2 - 2 \times 10 \times 14 \cos 43.3 (= 92.2)$ Evaluating square root  $(QR =) 9.6(0) (9.60 \text{ to } 9.603\dots)$ c.a.o. ww2	<b>M2</b> <b>M1</b>  <b>A1</b>	If <b>M0, M1</b> for correct implicit statement <b>M1 (dependent on M2)</b> for square root of correct combination (not negative) i.e $16 \cos 43.3 (11.64\dots)$ implies <b>M2M0</b> <span style="float: right;"><b>[7]</b></span>

<b>4 (a)</b>	$(AB =) \frac{250}{\sin 126} \times \sin 23$ (s.o.i by 120...) <b>121 (120.7 to 121) (m)</b> c.a.o. www3	<b>M2</b>  <b>A1</b>	<b>M1</b> for $\frac{AB}{\sin 23} = \frac{250}{\sin 126}$ o.e. (implicit)
<b>(b) (i)</b>	<b>280</b>	<b>B1</b>	
<b>(ii)</b>	<b>(0)69</b> c.a.o.	<b>B2</b>	<b>SC1</b> for answer 249 <span style="float: right;"><b>[6]</b></span>

<b>5 (a) (i)</b>	<b>1.5, 3.75, -1.5</b>	<b>B1, B1, B1</b>	
<b>(ii)</b>	12 points plotted ft Curve through at least 10 points and correct shape over full domain Two separate branches, one on each side of y-axis, neither in contact with y-axis	<b>P3 ft</b>  <b>C1</b>  <b>B1</b>	<b>P2 ft</b> for 10 or 11 points, <b>P1 ft</b> for 8 or 9 points i.s.w. if two branches joined  Independent
<b>(b)</b>	<b><math>-1.4 \leq x \leq -1.1</math> and <math>3.1 \leq x \leq 3.4</math></b>	<b>B1, B1</b>	i.s.w. 3rd answer if curve cuts $y = 1$ again
<b>(c) (i)</b>	Correct ruled tangent at $x = 2$ or $x = -2$ Evidence of rise/run  <b>0.8 to 1.2</b>	<b>M1</b> <b>M1</b>  <b>A1</b>	Long enough to be able to find gradient <b>Dependent</b> – check their graph against gradient of 1 – must be correct side of 1 <b>No tangent drawn M0M0</b>
<b>(ii)</b>	<b>0.8 to 1.2 inc.</b> or same answer as <b>(i)</b> ft	<b>B1 ft</b>	
<b>(d) (i)</b>	Correct ruled line to cut curve for <b>all</b> possible intersections (at least 2)	<b>B1</b>	Within $\frac{1}{2}$ square of $(-1, 1)$ and $(1, -1)$
<b>(ii)</b>	<b>-1.3 to -1.05, 1.05 to 1.3</b> inclusive	<b>B1, B1</b>	i.s.w. any extra answers
<b>(e)</b>	$y = kx$ with $k \geq \frac{1}{2}$ o.e. or $x = 0$	<b>B2</b>	If <b>B0</b> , allow <b>SC1</b> for $y = kx$ with $k < \frac{1}{2}$ or for y-axis stated <span style="float: right;"><b>[19]</b></span>

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2009	0580, 0581	04

<b>6 (a) (i)</b>	$0.5 [(x + 6) + (x + 2)] \times (x + 1) (= 40)$ or better $0.5(2x + 8)(x + 1) (= 40)$ o.e. $0.5(2x^2 + 10x + 8) (= 40)$ o.e. $x^2 + 5x + 4 = 40$ o.e. $x^2 + 5x - 36 = 0$	<b>M1A1</b>  <b>E1</b>	<b>M1</b> for any algebraic use of half base $\times$ height (Brackets may be implied later) May be first line If this first line, then <b>M0</b> <b>Dependent on M1A1.</b> Fully established – no errors throughout and at least 2 steps, one with 40 or 80, after first line
<b>(ii)</b>	<b>-9, 4</b>	<b>B1, B1</b>	<b>If B0, SC1</b> for +9 and -4
<b>(iii)</b>	$(BC^2 = )$ (their $x + 1$ ) <sup>2</sup> + (their $x + 2$ ) <sup>2</sup> $(BC = )$ <b>7.81(0...)</b> c.a.o. www2	<b>M1</b> <b>A1</b>	Their $x$ must be positive Ignore any extra solutions
<b>(b) (i)</b>	$9\frac{5}{12}$ or $\frac{108 + 5}{12}$ or $\frac{9 \times 12 + 5}{12}$ or $\frac{565}{60}$ or $\frac{9 \times 60 + 25}{60}$ seen	<b>E1</b>	Must be fractional form Condone $113/12 \times 60 = 565$ ; $9 \times 60 + 25 = 565$ Not for decimals
<b>(ii)</b>	$\frac{3y + 2}{3}$ or $\frac{y + 4}{2}$ o.e. $\frac{2(3y + 2)}{6} + \frac{3(y + 4)}{6}$ o.e.	<b>B1</b>  <b>B1</b>	or $\frac{6y + 4}{6} + \frac{3y + 12}{6}$ o.e.
<b>(iii)</b>	$\frac{2(9y + 16)}{12} = \frac{113}{12}$ o.e. $y = 4.5$ c.a.o. www2	<b>M1</b> <b>A1</b>	o.e. means with common denominator or better (Trial and error scores 2 or 0.)
<b>(iv)</b>	(Total dist =) $(3 \times \text{their } y) + 2 + (\text{their } y) + 4$ o.e. (Average speed =) $\frac{\text{their } 24}{9\frac{5}{12}}$ o.e. <b>2.55</b> (km/h) (2.548 – 2.549) c.a.o. www 3	<b>M1</b>  <b>M1</b> <b>A1</b>	(= 24) (dependent) Must be km divided by hours o.e. for full method Accept fractions in range

[15]

<b>Page 5</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – May/June 2009</b>	<b>0580, 0581</b>	<b>04</b>

<b>7 (a)</b>	$250x^2 = 4840$ o.e. $x^2 = 19.36$ or $(x =) \sqrt{4840 \div 250} (= 4.4)$	<b>M1</b> <b>E1</b>	Allow <b>M1</b> for $250 \times 4.4^2 = 4840$ Then <b>E1</b> for $250 \times 19.36 = 4840$
<b>(b)</b>	<b>42.6</b> (kg) cao (42.592 or 42.59)	<b>B2</b>	<b>SC1</b> for figures 426 or 4259...
<b>(c)</b>	<b>26.4</b> (cm) c.a.o.	<b>B2</b>	If <b>B0, M1</b> for any of following $88 \div 4.4 = 20$ and $120 \div 20 = 6$ (accept 6 bars high o.e.) or $88h = 4.4^2 \times 120$ or $250 \times 88 \times h = 120 \times 4840$
<b>(d) (i)</b>	$4840 \div 4200$ (implied by 1.15(2)) $\div \frac{4}{3}\pi$ (implied by 0.274 to 0.276) $\sqrt[3]{\quad}$ (seen or implied by correct answer to more than 2 dp) <b>0.649 – 0.651</b>	<b>M1</b> <b>M1</b> <b>M1 dep</b> <b>A1</b>	$4200 \times \frac{4}{3}\pi r^3 = 4840$ $(r^3 =) 4840 \div (4200 \times \frac{4}{3}\pi)$ $\sqrt[3]{\quad}$ Third M <b>dependent</b> on <b>M1M1</b> Must be 3dp or better
<b>(ii)</b>	<b>5.31 (5.306 – 5.31)</b> (cm <sup>2</sup> )	<b>B1</b>	
<b>(iii)</b>	$\frac{4200 \times \text{their (ii)}}{2 \times 4.4^2 + 4 \times 4.4 \times 250} \times 100$ <b>501.9 – 503</b> (%) c.a.o. www4	<b>M3</b>  <b>A1</b>	If <b>M0, M1</b> for $4200 \times \text{their (ii)}$ (22299) <b>and M1</b> (independent) for correct method for surface area of solid cuboid (4438.72)  <b>[15]</b>

<b>8</b>			Throughout the question ratios score zero. If using decimals, 2 s.f. correct answers to parts <b>(c)</b> and <b>(d)</b> – penalty of 1 once Use of words e.g. 1 in 400 or 1 out of 400, Correct answers – penalty of one For method marks only accept probabilities $p$ and $q$ between 0 and 1
<b>(a)</b>	$p = \frac{1}{20}, q = \frac{19}{20}$ o.e.	<b>B1</b>	Could be on diagram
<b>(b) (i)</b>	$\frac{1}{400}$ o.e. c.a.o.	<b>B2</b>	0.0025 allow <b>M1</b> for $(\text{their } p)^2$ o.e.
<b>(ii)</b>	$\frac{38}{400}$ o.e. c.a.o.	<b>B2</b>	0.095 allow <b>M1</b> for 2 (their $p$ )(their $q$ ) o.e.
<b>(c)</b>	$\frac{38}{8000}$ o.e. c.a.o.	<b>B2</b>	0.00475 allow <b>M1</b> for 2(their $p$ ) <sup>2</sup> (their $q$ ) o.e. including their <b>(ii)</b> $\times$ their $p$
<b>(d)</b>	their <b>(b)(i)</b> + their <b>(c)</b> $\frac{58}{8000}$ o.e. c.a.o.	<b>M1</b> <b>A1</b>	0.00725
<b>(e)</b>	their <b>(d)</b> $\times 1000 = 7.25$ o.e. ft	<b>B1 ft</b>	Accept 7 or 8 or an equivalent integer ft <b>[10]</b>

<b>Page 6</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – May/June 2009</b>	<b>0580, 0581</b>	<b>04</b>

<b>9 (a) (i)</b>	<b>174 to 174.25</b> (cm) c.a.o.	<b>B1</b>	
<b>(ii)</b>	<b>167</b> (cm) c.a.o.	<b>B1</b>	
<b>(iii)</b>	<b>12</b> (cm) c.a.o.	<b>B1</b>	
<b>(iv)</b>	<b>37</b> c.a.o.	<b>B2</b>	If <b>B0, B1</b> for 63 seen in working space
<b>(b) (i)</b>	<b>10, 25</b>	<b>B1</b>	s.o.i. allow 1 slip Use of $\Sigma fx$ where the $x$ 's are in/on their intervals (allow one more slip) (17 230)  (dependent on second M) $\div 100$
<b>(ii)</b>	155, 165, 175, 185 (their $10 \times 155 + \text{their } 25 \times 165 + 47 \times 175 + 18 \times 185$ )	<b>M1</b>	
	$\div 100$	<b>M1</b>	
	<b>172 or 172.3</b> (cm) c.a.o. www 4	<b>A1</b>	

<b>10 (a) (i)</b>	<b>-2,</b>	<b>B1</b>	
<b>(ii)</b>	<b>26,</b>	<b>B1</b>	
<b>(iii)</b>	$\frac{1}{8}$ o.e.	<b>B1</b>	
<b>(b)</b>	$\frac{y+1}{2} (= x)$ ( $f^{-1}(x) = $ ) $\frac{x+1}{2}$ o.e. www2	<b>M1</b> <b>A1</b>	If switch $x$ and $y$ first then <b>M1</b> for $x = 2y - 1$ or If use a diagram/chart then <b>M1</b> for any evidence of $+1$ then result $\div 2$
<b>(c)</b>	$z = x^2 + 1$ $z - 1 = x^2$  ( $x = $ ) $\sqrt{z-1}$ www2	<b>M1</b> <b>M1</b>	Correct rearrangement at any stage for $x$ or $x^2$ . Correct sq root at any stage Ignore $+$ , $-$ or $\pm$ in front of $\sqrt{\quad}$
<b>(d)</b>	$(2x-1)^2 + 1$ $= 4x^2 - 4x + 2$ or $2(2x^2 - 2x + 1)$ www 2	<b>M1</b> <b>A1</b>	Final answer but condone one minor factorising slip if first answer seen
<b>(e)</b>	9	<b>B1</b>	
<b>(f)</b>	$2(2x-1) + x^2 + 1 (= 0)$ or better ( $x^2 + 4x - 1 = 0$ )  ( $x = $ ) $\frac{-4 \pm \sqrt{4^2 - 4(1)(-1)}}{2 \times 1}$ ft  ( $x = $ ) <b>-4.24, 0.24</b> c.a.o. www 4 (final answers)	<b>B1</b> <b>M1</b> <b>M1</b>  <b>A1,A1</b>	$\sqrt{4^2 - 4(1)(-1)}$ or better seen If in form $\frac{p \text{ or } -\sqrt{q}}{r}$ for $-4$ and $2 \times 1$ or better <b>Ft</b> their 1, 4 and $-1$ from quadratic equation seen <b>After A0A0, SC1</b> for $-4.2$ or $-4.235$ or $-4.236\dots$ <b>and</b> $0.2$ or $0.235$ or $0.236\dots$ The <b>SC1</b> 's www imply the <b>M</b> marks
<b>(g) (i)</b>	Straight line with positive gradient and negative $y$ -intercept	<b>L1</b>	
<b>(ii)</b>	U-shape Parabola vertex on positive $y$ -axis	<b>C1</b> <b>V1</b>	
			<b>Dependent</b> <b>[18]</b>

<b>11 (a)</b>	<b>15, 21, 28, 36</b>	<b>B2</b>	<b>B1</b> for 3 correct
<b>(b) (i)</b>	$10 + 15 = 25, 15 + 21 = 36$ etc	<b>B1</b>	Any two complete and correct statements
<b>(ii)</b>	<b>Square</b>	<b>B1</b>	
<b>(c) (i)</b>	<b>2</b>	<b>B1</b>	
<b>(ii)</b>	$\frac{4 \times 5}{2} = 10$ o.e.	<b>E1</b>	
<b>(iii)</b>	<b>16 290</b> c.a.o.	<b>B1</b>	
<b>(d) (i)</b>	$\frac{(n+1)(n+2)}{2}$ or $\frac{n^2 + 3n + 2}{2}$ seen $\frac{n(n+1)}{2} + \frac{(n+1)(n+2)}{2}$ or $\frac{n^2 + n}{2} + \frac{n^2 + 3n + 2}{2}$ $\frac{(n+1)(n+n+2)}{2}$ $\frac{2n^2 + 4n + 2}{2}$ $\frac{(n+1)(2n+2)}{2}$ $n^2 + 2n + 1$ $\frac{2(n+1)(n+1)}{2} = (n+1)^2$ $(n+1)^2$	<b>M1</b> <b>M1</b> <b>E1</b>	Denominator could be their $k$ May be implied by next line This line must be seen and at least one more step, without any error, to gain the E mark <b>Dependent on M1M1.</b> Fully established – no errors
<b>(ii)</b>	<b>1711 and 1770</b> final answers c.a.o.	<b>B2</b>	<b>SC1</b> for <b>59 or 58 or 1711 or 1770</b> seen <b>[12]</b>

Graph for Question 5

